

## Tilburg University

### On measures of nonnormality of matrices

Elsner, L.; Paardekooper, M.H.C.

*Publication date:*  
1984

[Link to publication in Tilburg University Research Portal](#)

*Citation for published version (APA):*

Elsner, L., & Paardekooper, M. H. C. (1984). *On measures of nonnormality of matrices*. (Research memorandum / Tilburg University, Department of Economics; Vol. FEW 167). Unknown Publisher.

#### General rights

Copyright and moral rights for the publications made accessible in the public portal are retained by the authors and/or other copyright owners and it is a condition of accessing publications that users recognise and abide by the legal requirements associated with these rights.

- Users may download and print one copy of any publication from the public portal for the purpose of private study or research.
- You may not further distribute the material or use it for any profit-making activity or commercial gain
- You may freely distribute the URL identifying the publication in the public portal

#### Take down policy

If you believe that this document breaches copyright please contact us providing details, and we will remove access to the work immediately and investigate your claim.

CBM  
R

7626

1984

167



Bestemming

ECO


TIJDSCHRIFTENBUREAU  
BIBLIOTHEEK  
KATHOLIEKE  
HOGESCHOOL  
TILBURG

Nr.

941

subfaculteit der econometrie

## RESEARCH MEMORANDUM




\* C I N O O 4 1 0 \*

TILBURG UNIVERSITY

DEPARTMENT OF ECONOMICS

Postbus 90153 - 5000 LE Tilburg  
Netherlands



ON MEASURES OF NONNORMALITY OF MATRICES

L. Elsner, M.H.C. Paardekooper

December 1984

## ON MEASURES OF NONNORMALITY OF MATRICES

L. Elsner, Fakultät für Mathematik, Universität Bielefeld, Postfach  
8640, 4800 Bielefeld, FRG.

M.H.C. Paardekooper, Department of Econometrics, Tilburg University,  
5000 LE Tilburg, The Netherlands.

Abstract

This paper discusses measures  $\mu_i$ ,  $i = 1, \dots, 9$ , of nonnormality of matrices and their interrelations. Some of the measures are new. Each non-negative measure  $\mu_i(A)$  equals zero iff  $A$  is normal. We compare these measures with several new inequalities. Some of these comparisons, e.g., (C12), (C15) and (C16), manifest wellknown phenomena of ill conditioned eigenproblems.

1. Introduction

The class of nonnormal matrices has received some attention of numerical analysts. In particular in connection with certain eigenvalue algorithms normal and nonnormal matrices show quite different behaviour. Related to this fact is the difference in the sensitivity of the eigenvalues and eigenvectors under perturbations of the entries of the matrix.

For analyzing these difficulties several measures of nonnormality have appeared in the literature. We give here an overview of the measures used, introduce some new ones and give in particular comparisons between them.

Throughout this paper  $n > 1$  is a fixed integer. Let  $C^{n,n}$  denote the set of all complex  $n \times n$  matrices and  $A \in C^{n,n}$ . We associate with  $A$  the following magnitudes.

- its eigenvalues  $\lambda_i$  and its singular values  $\sigma_i$ , ordered such that



$$|\lambda_1| \geq |\lambda_2| \geq \dots \geq |\lambda_n|, \quad \sigma_1 \geq \sigma_2 \geq \dots \geq \sigma_n;$$

- its polar factors  $H_1, H_2$ , i.e., the uniquely determined positive semi-definite square roots of  $AA^*$  and  $A^*A$ ;
- $\alpha_i$ ,  $i = 1, \dots, n$ , with  $\alpha_1 \geq \alpha_2 \geq \dots \geq \alpha_n$ , the eigenvalues of the hermitean part  $(A+A^*)$  of  $A$ ;
- $\beta_i$ ,  $i = 1, \dots, n$ , with  $\beta_1 \geq \beta_2 \geq \dots \geq \beta_n$ , the eigenvalues of the skew-hermitean part  $(A-A^*)/(2i)$  of  $A$ .

$A$  is normal, if  $A^*A - AA^* = 0$  holds. Let  $N$  denote the set of all normal matrices in  $C^{n,n}$ ,  $U$  the set of all unitary matrices in  $C^{n,n}$ , and  $D$  the set of diagonal matrices in  $C^{n,n}$ . We denote by  $\|\cdot\|_2$  the spectral norm and by  $\|\cdot\|_F$  the Frobenius norm of a matrix. For  $X$  nonsingular  $\kappa_1(X) := \|X\|_1 \|X^{-1}\|_1$ ,  $i = 2, F$  is the condition number of  $X$ .

## 2. Characterization of normal matrices

There are quite a few characterizations for  $A$  being normal.

**Theorem 1.** For  $A \in C^{n,n}$  with eigenvalues  $\{\lambda_i\}$ , singular values  $\{\sigma_i\}$ , polar factors  $H_i$  ( $i = 1, 2$ ) and eigenvalues  $\{\alpha_i\}$ ,  $\{\beta_i\}$  of  $(A+A^*)/2$  and  $(A-A^*)/(2i)$  resp. as above the following are equivalent.

- (i)  $A$  is normal, i.e.,  $A^*A - AA^* = 0$ ;
- (ii) There exists a  $V \in U$  such that  $V^*AV = \text{diag}(\lambda_i)$ ;
- (iii)  $\|A\|_F^2 - \sum_{i=1}^n |\lambda_i|^2 = 0$ ;
- (iv)  $A = X \text{diag}(\lambda_i) X^{-1}$  for some  $X \in C^{n,n}$  nonsingular and  $\kappa_2(X) = 1$ ;
- (v)  $|\lambda_i| - \sigma_i = 0$ ,  $i = 1, \dots, n$ ;
- (vi)  $H_1 = H_2$ ;
- (vii) There exists a  $V \in U$  such that both  $V^* \frac{A+A^*}{2} V$  and  $V^* \frac{A-A^*}{2} V$  are diagonal;
- (viii) There exist permutations  $p$  and  $q$  of  $\{1, \dots, n\}$  such that

$$\lambda_j = \alpha_{p(j)} + i \beta_{q(j)}, \quad j = 1, \dots, n. \quad \square$$

We refrain from giving a proof, as the results are either trivial or easy consequences of the quantitative statements in theorem 2.

We make one exception. (viii) implies (iii) as can be seen from the simple equalities

$$\sum_{j=1}^n (\alpha_{p(j)}^2 + \beta_{q(j)}^2) = \left\| \frac{A+A^*}{2} \right\|_F^2 + \left\| \frac{A-A^*}{2i} \right\|_F^2 = \|A\|_F^2.$$

### 3. Measures of Nonnormality of Matrices

Theorem 1 motivates the introduction of several measures of nonnormality.

The most natural measure of nonnormality seems to be

$$\mu_1(A) := \min \{ \|A - N\|_F \mid N \in N \},$$

and

$$\tilde{\mu}_1(A) := \min \{ \|A - N\|_2 \mid N \in N \}.$$

Besides this we consider

$$\mu_2(A) := \|A^*A - AA^*\|_F, \quad \tilde{\mu}_2(A) := \|A^*A - AA^*\|_2,$$

$$\mu_3(A) := (\|A\|_F^2 - \sum_{i=1}^n |\lambda_i|^2)^{\frac{1}{2}},$$

$$\mu_4(A) := \max_i |\lambda_i - \sigma_i|,$$

$$\mu_5(A) := \|H_1 - H_2\|_F,$$

$$\mu_6(A) := \min \left\{ \left( \sum_{j=1}^n |\lambda_j - (\alpha_{p(j)} + i\beta_{q(j)})|^2 \right)^{\frac{1}{2}}, p, q \text{ permute } \{1, \dots, n\} \right\}$$

$$\mu_7(A) := \min \{ \|U - V\|_F \mid U^*(A + A^*)U, V^*(A - A^*)V \in D, U, V \in U \},$$

$$\tilde{\mu}_7(A) := \min \{ \|U - V\|_2 \mid U^*(A + A^*)U, V^*(A - A^*)V \in D, U, V \in U \},$$

$$\mu_8(A) := \min \{ \|U - V\|_F \mid U A V^* \in D, U, V \in U \},$$

$$\tilde{\mu}_8(A) := \min \{ \|U - V\|_2 \mid U A V^* \in D, U, V \in U \},$$

and for  $A$  diagonalizable

$$\mu_9(A) := \min \{ \kappa_F(X) - n \mid X \in C^{n,n}, X^{-1} A X = \text{diag}(\lambda_1) = \Lambda \},$$

$$\hat{\mu}_9(A) := \min \{ \kappa_2(X) - 1 \mid X \in C^{n,n}, X^{-1} A X = \text{diag}(\lambda_1) = \Lambda \}.$$

It is easy to see that each of these measures is invariant with respect to unitary transformation, and conjugate transposition, i.e.,

$$\mu_i(A) = \mu_i(U^* A U), \quad U \text{ unitary}, \quad i = 1, \dots, 9,$$

$$\mu_i(A) = \mu_i(A^*),$$

and similarly for  $\hat{\mu}_i(A)$ ,  $i = 1, 2, 7, 8, 9$ .

These measures, except for  $\mu_4$ ,  $\mu_5$  and  $\mu_8$ ,  $\hat{\mu}_8$ , are also invariant with respect to shift transformation:

$$\mu_i(A) = \mu_i(A + wI), \quad w \in \mathbb{C}.$$

#### 4. Comparisons between Measures of Nonnormality

Comparisons between the measures  $\mu_i(A)$ ,  $\hat{\mu}_i(A)$  are given in Theorem 2. For  $A \in C^{n,n}$  the following singularities hold.

$$(C0) \quad \hat{\mu}_i \leq \mu_i \leq \sqrt{n} \hat{\mu}_i, \quad i = 1, 2, 7, 8;$$

$$(C1) \quad \frac{\mu_2^2}{\sqrt{6} \|A\|_F} \leq \mu_3 \leq \left( \frac{n^3 - n}{12} \right)^{\frac{1}{2}} \mu_2;$$

$$(C2) \quad \mu_2^2 \leq \sqrt{2 \left( \|A\|_F^2 + \sum_{i=1}^n \lambda_i^2 \right)} \leq 2 \|A\|_F \mu_3;$$

$$(C3) \quad \mu_2^2 \leq 4 \|A\|_2 \mu_1;$$

$$(C4) \quad \mu_2^2 \leq 4 \|A\|_F \hat{\mu}_1 + 2\sqrt{n} \hat{\mu}_1^2;$$



$$(C5) \quad \mu_1 \leq \mu_3 ;$$

$$(C6) \quad \mu_4 \leq \mu_3 ;$$

$$(C7) \quad \mu_3^2 \leq 2\sqrt{n} \|A\|_F \mu_4 \leq 2n \|A\|_2 \mu_4 ;$$

$$(C8) \quad \|A\|_2^{+1} \mu_5 \leq \mu_2^2 \leq 2 \|A\|_2 \mu_5 ;$$

$$(C9) \quad \tilde{\mu}_2^2 \leq 2 \|A\|_2^2 \tilde{\mu}_8 ;$$

$$(C10) \quad \tilde{\mu}_2^2 \leq 8 \|A\|_2^2 \tilde{\mu}_7 ;$$

$$(C11) \quad \mu_6^2 \leq \mu_3^2 \leq \max\{\|A\|_F \mu_6, 2 \|A\|_F \mu_6 - \mu_6^2\} \leq 2 \|A\|_F \mu_6 ;$$

$$(C12) \quad \tilde{\mu}_9^2 (1+\tilde{\mu}_9)^{-1} \leq \mu_9 \leq \frac{1}{2} n \tilde{\mu}_9 (1+\tilde{\mu}_9)^{-1},$$

and if all eigenvalues  $\lambda_j$  of  $A$  are simple, then

$$\mu_9 \leq \sum_{j=1}^n (1+(n-1)^{-1} \delta_j^{-2} \mu_3^2)^{\frac{n-1}{2}} - n ,$$

where  $\delta_j = \min\{|\lambda_j - \lambda_1| \mid i \neq j\}$ ;

$$(C13) \quad \mu_2^2 \leq 2 \|A\|_2 \|A\|_F \tilde{\mu}_9 (2+\tilde{\mu}_9) \leq 2 \|A\|_F^2 \tilde{\mu}_9 (2+\tilde{\mu}_9),$$

$$(C14) \quad \mu_3^2 \leq \|A\|_F^2 \tilde{\mu}_9 (2+\tilde{\mu}_9) \leq \|A\|_F^2 \tilde{\mu}_9 (2+\tilde{\mu}_9).$$

$$(C15) \quad \text{if } \mu_5 < \tau := \min\{|\sigma_1 - \sigma_j| \mid \sigma_1 \neq \sigma_j\}, \text{ then}$$

$$\mu_8 \leq (n+1) \tau^{-1} \mu_5;$$

$$(C16) \quad \text{if the eigenvalues } \beta_j, j = 1, \dots, n, \text{ of } (A-A^*)/(2i) \text{ are pairwise distinct and}$$

$$\mu_2^2 < (6-4\sqrt{2}) \delta \tilde{\delta},$$

where

then

$$\mu_7 \leq \sqrt{n}(\delta\hat{\delta} - \frac{3}{2} \mu_2^2)^{-1} \mu_2^2.$$

In all these comparisons

$$\mu_i = \mu_i(A), i = 1, \dots, 9 \text{ and } \hat{\mu}_1 = \mu_1(A), i = 1, 2, 7, 8, 9.$$

The comparisons in this theorem can be visualized with a directed graph, see fig. 1. There is a directed edge from node  $\mu_i$  (, or  $\hat{\mu}_i$ ) to  $\mu_j$  (, or  $\hat{\mu}_j$ ),  $i \neq j$ , iff

$$(4.1) \quad \mu_i(A) \leq \varphi(\mu_j(A))$$

for some monotonically continuous  $\varphi$  with  $\lim_{t \rightarrow 0} \varphi(t) = 0$ .




Fig. 1. The directed graph for equivalent measures.

The measures  $\mu_i$ ,  $i = 1, \dots, 9$  and  $\hat{\mu}_i$ ,  $i = 1, 2, 7, 8, 9$  are nodes in a completely connected graph, for relation 4.1 is transitive.

Proof of theorem 2. Assume  $\hat{\mu}_1 = \|A-\hat{N}\|_2$ ,  $\mu_1 = \|A-\hat{N}\|_F$ ,  $\hat{N}, \hat{N} \in \mathcal{N}$ . Then

$$\hat{\mu}_1 = \|A-\hat{N}\|_2 \leq \|A-\hat{N}\|_2 \leq \|A-\hat{N}\|_F = \mu_1$$

and

$$\mu_1 = \|A-\hat{N}\|_F \leq \|A-\hat{N}\|_F \leq \sqrt{n} \|A-\hat{N}\|_2 = \sqrt{n} \hat{\mu}_1.$$

The same reasoning applies to  $\mu_i$ ,  $\hat{\mu}_i$ ,  $i = 2, 7, 8$ . This proves (C0).

The first inequality in (C1) is a result of Eberlein [1], the second is a result of Henrici [2] in its original form.

The second inequality of (C2) is a consequence of

$$\sum_{i=1}^n |\lambda_i|^2 \leq \|A\|_F^2 = \sum_{i=1}^n \sigma_i^2$$

(also known as "Schurs Lemma"), the first inequality is just a rearrangement of the inequality

$$(4.2) \quad \left( \sum_{i=1}^n |\lambda_i|^2 \right) \leq \|A\|_F^4 - \frac{1}{2} \mu_2^4,$$

established by Kress-de Vries-Wegmann [3].

For the proof of (C3) we use that for  $N \in \mathcal{N}$

$$(4.3) \quad A^* A - A A^* = A^* (A-N) - (A-N) N^* + (A-N)^* N - A (A-N)^*$$

holds. Hence for any  $N \in \mathcal{N}$  (, with the wellknown inequality  $\|AB\|_F \leq \|A\|_2 \|B\|_F$ ),

$$(4.4) \quad \mu_2^2 \leq 2(\|A\|_2 + \|N\|_2) \|A-N\|_F.$$

If  $N$  is such that  $\mu_1 = \|A-N\|_F$  and  $U^* N U = D \in \mathcal{D}$  for  $U \in \mathcal{U}$ , then it is obvious from

$$(4.5) \quad \mu_1 = \|A-N\|_F = \min \{ \|U^*AU-D\|_F \mid D \in D, U \in U \}$$

that  $D$  is the diagonal part of  $U^*AU$ . In particular

$$(4.6) \quad \|N\|_2 = \|D\|_2 \leq \|A\|_2.$$

As follows from the definition of the spectral norm (4.4) and (4.6) together yield (C3).

Similarly from (4.3)

$$(4.7) \quad \mu_2^2 \leq 2(\|A\|_F + \|N\|_F) \|A-N\|_2,$$

and, using in (4.7)

$$\|N\|_F \leq \|A-N\|_F + \|A\|_F \leq \sqrt{n} \|A-N\|_2 + \|A\|_F,$$

(C4) follows.

For (C5) we consider a unitary matrix  $V$  that transforms  $A$  to upper triangular form:

$$(4.8) \quad V^*AV = \Lambda + M.$$

Then

$$\|A\|_F^2 = \|\Lambda\|_F^2 + \|M\|_F^2.$$

Hence

$$\mu_3 = \|M\|_F = \|A-V^*\Lambda V\|_F.$$

This implies  $\mu_3 \geq \mu_1$ .

(C6) is a result of Ruhe [5], while its counterpart (C7) is an easy consequence of

$$\mu_3^2 = \sum_{i=1}^n (\sigma_i^2 - |\lambda_i|^2) = \sum_{i=1}^n (\sigma_i + |\lambda_i|)(\sigma_i - |\lambda_i|)$$

$$\begin{aligned}
&\leq \mu_4 \sum_{i=1}^n (\sigma_i + |\lambda_i|) \leq \mu_4 (\sqrt{n} (\sum_{i=1}^n \sigma_i^2)^{\frac{1}{2}} + \sqrt{n} (\sum_{i=1}^n |\lambda_i|^2)^{\frac{1}{2}}) \\
&\leq 2 \sqrt{n} \|A\|_F \mu_4.
\end{aligned}$$

The second inequality in (C7) follows from  $\|B\|_F \leq \sqrt{n} \|B\|_2$  for every  $B \in \mathbb{C}^{n,n}$ .

For the proof of the second inequality of (C8) we make use of the singular value decomposition

$$(4.9) \quad A = W \Sigma V^*,$$

where  $W, V \in U$  and  $\Sigma = \text{diag}(\sigma_i)$ . As

$$A = W \Sigma W^* W V^*,$$

and

$$A = W V^* V \Sigma V^*,$$

we have that

$$A = H_1 U = U H_2,$$

where

$$H_1 = W \Sigma W^*, H_2 = V \Sigma V^* \text{ and } U = W V^*.$$

An easy calculation gives

$$(4.10) \quad H_1 - H_2 = W(\Sigma Y - Y \Sigma) V^*$$

and

$$(4.11) \quad A^* A - A A^* = W(Y \Sigma^2 - \Sigma^2 Y) V^*$$

where

$$Y = W^* V = (y_{ij}) \in U.$$

Hence from (4.10)

$$(4.12) \quad \mu_5^2 = \|H_1 - H_2\|_F^2 = \|\Sigma Y - Y \Sigma\|_F^2 = \sum_{i,j=1}^n |y_{ij}|^2 (\sigma_i - \sigma_j)^2$$

where by (4.11)

$$(4.13) \quad \mu_2^4 = \|A^* A - A A^*\|_F^2 = \sum_{i,j=1}^n |y_{ij}|^2 (\sigma_i^2 - \sigma_j^2)^2 = \sum_{i,j=1}^n |y_{ij}|^2 (\sigma_i - \sigma_j)^2 (\sigma_i + \sigma_j)^2.$$

This shows

$$\mu_2^4 \leq 4 \sigma_1^2 \mu_5^2 = 4 \|A\|_2^2 \mu_5^2,$$

i.e., the second inequality of (C8).

To prove the first inequality of (C8) we observe that with

$$\|A^+\|_2 = \max\{\sigma_i^{-1} \mid \sigma_i > 0, i = 1, \dots, n\}$$

one has

$$|\sigma_i - \sigma_j| \leq \|A^+\|_2 |\sigma_i^2 - \sigma_j^2|.$$

(4.12) and (4.13) show now

$$\mu_5^2 \leq \|A^+\|_2^2 \mu_4^2$$

i.e., the first inequality of (C8).

For the proof of inequality (C9) we assume that  $\tilde{\mu}_8 = \|U_0 - V_0\|_2$ , where  $U_0, V_0 \in U$  and  $U_0^* A V_0^* = D \in D$ . Then

$$A^* A - A A^* = V_0^* D^* D V_0 - U_0^* D^* D U_0$$



$$= (V_0 - U_0)^* D^* D V_0 - U_0^* D^* D (V_0 - U_0).$$

Hence

$$\tilde{\mu}_2^2 \leq 2 \|D\|_2^2 \|V_0 - U_0\|_2^2 = 2 \|A\|_2^2 \tilde{\mu}_8.$$

For the proof of (C10) we assume that  $\tilde{\mu}_7 = \|U_1 - V_1\|_2$  where

$$U_1^* \frac{A+A^*}{2} U_1 = M \in D, \quad V_1^* \frac{A-A^*}{2} V_1 = N \in D,$$

with

$$U_1, V_1 \in U.$$

So

$$A = U_1 M U_1^* + V_1 N V_1^*, \quad A^* = U_1 M U_1^* - V_1 N V_1^*.$$

Let be

$$R = V_1 - U_1.$$

Then

$$\frac{1}{2} (A^* A - A A^*) = U_1 M U_1 V_1^* N V_1^* - V_1 N V_1^* U_1 M U_1^*.$$

Hence, with  $W = V_1^* U_1$ ,

$$\begin{aligned} V_1^* \cdot \frac{1}{2} (A^* A - A A^*) V_1 &= W M W^* N - N W M W^* \\ &= (W - I) M W^* N + M (W^* - I) N - N M (W^* - I) - N (W - I) M W^*. \end{aligned}$$

Consequently,

$$\frac{1}{2} \tilde{\mu}_2^2 \leq \| (W - I) M W^* N \|_2^2 + \| M (W^* - I) N \|_2^2 + \| N M (W^* - I) \|_2^2 + \| N (W - I) M W^* \|_2^2$$

$$\leq 4 \|W-I\|_2 \|N\|_2 \|M\|_2 \leq 4 \|U_1 - V_1\|_2 \|A\|_2^2 = 4 \|A\|_2^2 \tilde{\mu}_7.$$

This proves (C10).

Without loss of generality we may assume in the proof of the first inequality of (C11) that  $A$  is an uppertriangular matrix since both  $\mu_6$  as  $\mu_3$  are unitary invariants of  $A$ :  $\mu_6(A) = \mu_6(U^*AU)$ ,  $\mu_3(U^*AU) = \mu_3(A)$  for  $U \in \mathcal{U}$ . So let be

$$A = \Lambda + R,$$

where

$$r_{ij} = 0, \quad 1 \leq j < i \leq n$$

and

$$\Lambda = \text{diag}(\lambda_j),$$

with

$$\lambda_j = \tau_j + i \nu_j, \quad j = 1, \dots, n.$$

Then

$$\left\| \frac{A+A^*}{2} - \frac{\Lambda+\Lambda^*}{2} \right\|_F^2 = \left\| \frac{1}{2}(R+R^*) \right\|_F^2 = \frac{1}{2} \|R\|_F^2 = \frac{1}{2} \mu_3^2.$$

According to the Hoffman-Wielandt theorem [8] there exists a permutation  $j \rightarrow p(j)$  of  $\{1, \dots, n\}$  such that

$$\sum_{j=1}^n (\tau_j - \alpha_{p(j)})^2 \leq \left\| \frac{A+A^*}{2} - \frac{\Lambda+\Lambda^*}{2} \right\|_F^2 = \frac{1}{2} \mu_3^2.$$

Similarly one finds for the skewhermitean part

$$\left\| \frac{A-A^*}{2} - \frac{\Lambda-\Lambda^*}{2} \right\|_F^2 = \left\| \frac{1}{2}(R-R^*) \right\|_F^2 = \frac{1}{2} \|R\|_F^2 = \frac{1}{2} \mu_3^2.$$

For some permutation  $j \rightarrow q(j)$  of  $\{1, \dots, n\}$

$$\sum_{j=1}^n (v_j - \beta_{q(j)})^2 \leq \frac{1}{2} \mu_3^2.$$

Consequently

$$\mu_6 \leq \mu_3.$$

So far as concerns the second inequality of (C11) observe

$$\begin{aligned} \mu_3^2 &= \|A\|_F^2 - \sum_{j=1}^n |\lambda_j|^2 = \left\| \frac{A+A^*}{2} \right\|_F^2 + \left\| \frac{A-A^*}{2} \right\|^2 - \sum_{j=1}^n |\lambda_j|^2 \\ &= \sum_{j=1}^n (\alpha_j^2 + \beta_j^2 - |\lambda_j|^2). \end{aligned}$$

Assume  $\lambda_j = \tau_j + i v_j$ ,  $j = 1, \dots, n$ . Then

$$\mu_3^2 = \sum_{j=1}^n (\alpha_{p(j)}^2 - \tau_j^2) + \sum_{j=1}^n (\beta_{q(j)}^2 - v_j^2),$$

where the permutations  $p, q$  realize the minimum that occur in the definition of  $\mu_6$ . Hence

$$\begin{aligned} \mu_3^2 &= \sum_{j=1}^n (\alpha_{p(j)} - \tau_j)(\alpha_{p(j)} + \tau_j) + \sum_{j=1}^n (\beta_{q(j)} - v_j)(\beta_{q(j)} + v_j) \\ &\leq \mu_6 \left( \sum_{j=1}^n ((\alpha_{p(j)} + \tau_j)^2 + (\beta_{q(j)} + v_j)^2) \right)^{\frac{1}{2}} \\ &\leq \mu_6 \left( \left( \sum_{j=1}^n (\alpha_j^2 + \beta_j^2) \right)^{\frac{1}{2}} + \left( \sum_{j=1}^n (\tau_j^2 + v_j^2) \right)^{\frac{1}{2}} \right) \\ &= \mu_6 (\|A\|_F + \left( \sum_{j=1}^n |\lambda_j|^2 \right)^{\frac{1}{2}}) = \mu_6 (\|A\|_F + (\|A\|_F^2 - \mu_3^2)^{\frac{1}{2}}). \end{aligned}$$

The last equation gives

$$\mu_3^2 - \|A\|_F \mu_6 \leq (\|A\|_F^2 - \mu_3^2)^{\frac{1}{2}} \mu_6.$$

Either

$$\mu_3^2 \leq \|A\|_F \mu_6,$$

or by squaring

$$\mu_3^2 \leq 2 \|A\|_F \mu_6 - \mu_6^2.$$

Hence

$$\mu_3^2 \leq \max \{ \|A\|_F \mu_6, 2 \|A\|_F \mu_6 - \mu_6^2 \} \leq 2 \|A\|_F \mu_6.$$

This proves (C11).

The results of (C12) are due to Smith [6]. There he proves, for

$$K := \inf \{ \kappa_F(X) \mid X^{-1} A X \in D, X \in C^{n,n} \}$$

and

$$k := \inf \{ \kappa_2(X) \mid X^{-1} A X \in D, X \in C^{n,n} \}$$

that

$$n - 2 + k + k^{-1} \leq K \leq \frac{1}{2} n (k + k^{-1}).$$

Since

$$\mu_9 = K - n, \quad \hat{\mu}_9 = k - 1$$

this yields

$$\hat{\mu}_9^2 (1 + \hat{\mu}_9)^{-1} \leq \mu_9 \leq \frac{1}{2} n \hat{\mu}_9 (1 + \hat{\mu}_9)^{-1}.$$

In the same paper Smith derives, if all eigenvalues of  $A$  are simple, with

$$\delta_j = \min_{i \neq j} |\lambda_j - \lambda_i|.$$

that

$$K \leq \sum_{j=1}^n (1+(n-1)^{-1} \delta_j^{-2} \mu_3^2)^{-1},$$

being the last inequality in (C12).

For the proof of (C13) we use the following facts:

a. If  $S$  is hermitean and  $X$  nonsingular, then Schurs Lemma applied to  $X^{-1} S X$  yields

$$(4.14) \quad \|S\|_F \leq \|X^{-1} S X\|_F.$$

b. If  $Y$  is positive definite and  $G \in \mathbb{C}^{n,n}$ , then

$$(4.15) \quad \|Y G Y^{-1} - G\|_F \leq (\kappa_2(Y)-1) \|G\|_F.$$

This can be shown by writing the linear operator

$$L : \mathbb{C}^{n,n} \rightarrow \mathbb{C}^{n,n}, \quad L(G) := Y G Y^{-1} - G$$

in the usual vectorized form (, see Marcus-Minc, [4], p. 9):

$\text{vec}(L(G)) = (Y^{-1} \otimes Y - I_n \otimes I_n) \text{vec}(G) = \hat{L} \text{vec}(G)$ , where  $\otimes$  denotes the Kronecker product. As  $\hat{L}$  is hermitean and has eigenvalues  $-1 + \eta_i / \eta_j$  ( $\eta_i$  eigenvalues of  $Y$ ) and as  $\|G\|_F$  is the usual euclidean norm of the vector  $\text{vec}(G) \in \mathbb{C}^{n^2}$ , the result (4.15) follows.

Assume  $\hat{\mu}_9 = \kappa_2(X_1)-1$ . From  $A = X_1^{-1} \Lambda X_1$  we derive the following relation

$$X_1 (A^* A - A A^*) X_1^{-1} = (Y \Lambda^* Y^{-1} - \Lambda^*) \Lambda - \Lambda (Y \Lambda^* Y^{-1} - \Lambda^*)$$

where

$$Y = X_1 X_1^* > 0.$$

Using (4.14) and (4.15) yields

$$\begin{aligned}
\mu_2^2 &\leq 2 \|\Lambda\|_2 \|\Lambda^* Y^{-1} - \Lambda^*\|_F \leq 2 \|\Lambda\|_2 \|\Lambda^*\|_F (\kappa_2(Y) - 1) \\
&= 2 \|\Lambda\|_2 \|\Lambda\|_F (\kappa_2^2(X_1) - 1) \\
&= 2 \|\Lambda\|_2 \|\Lambda\|_F \tilde{\mu}_9 (2 + \tilde{\mu}_9) \\
&\leq 2 \|\Lambda\|_F^2 \tilde{\mu}_9 (2 + \tilde{\mu}_9).
\end{aligned}$$

For the proof of (C14) we observe that  $A = X_1^{-1} \wedge X_1$  yields

$$\|\Lambda\|_F^2 \leq \kappa_2^2(X_1) \|\Lambda\|_F^2.$$

Hence, with  $\tilde{\mu}_9 = \kappa_2(X_1) - 1$ ,

$$\begin{aligned}
\mu_3^2 &= \|\Lambda\|_F^2 - \|\Lambda\|_F^2 \leq (\kappa_2^2(X_1) - 1) \|\Lambda\|_F^2 = \tilde{\mu}_9(2 + \tilde{\mu}_9) \|\Lambda\|_F^2 \\
&\leq \|\Lambda\|_F^2 \tilde{\mu}_9(2 + \tilde{\mu}_9),
\end{aligned}$$

i.e. (C14).

In the proof of inequality (C15) we make use of two lemmata. These describe the relation between several measures of non-unitarity of matrices. For that purpose we introduce the following notations

$$\pi_1(A) := \min\{\|A - U\|_F \mid U \in U\}, \quad \tilde{\pi}_1(A) := \{\|A - U\|_2 \mid U \in U\},$$

$$\pi_2(A) := \min\{\|UAU^* \Delta - I\|_F \mid U \in U, \Delta \in U \cap D\}, \quad \pi_3(A) := \|\Lambda^* A - I\|_F$$

$$\pi_2(A) := \min\{\|UAU^* \Delta - I\|_2 \mid U \in U, \Delta \in U \cap D\}, \quad \tilde{\pi}_3(A) := \|\Lambda^* A - I\|_2.$$

Lemma 1.  $\pi_1(A) = \pi_2(A)$ ,  $\tilde{\pi}_1(A) = \tilde{\pi}_2(A)$ .

Proof. Let be  $\pi_1(A) = \|A - V\|_F$ ,  $V \in U$ . Then there exists a unitary  $Q$  and a unitary diagonal matrix  $\Delta$  such that  $Q V Q^* \Delta = I$ . If  $A = V + E$ , then

$$\pi_2(A) \leq \|Q(V+E)Q^* \Delta - I\|_F = \|QE Q^* \Delta\|_F = \pi_1(A).$$



Reservely, let be  $\pi_2(A) = \|Q A Q^* \Delta^{-1}\|_F$  with  $Q \in U$  and  $\Delta \in U \cap D$ . If  $Q A Q^* \Delta = I + G$ , then  $A = Q^* (I+G) \Delta^* Q$  and  $Q^* \Delta^* Q \in U$ . So

$$\pi_1(A) \leq \|A - Q^* \Delta^* Q\|_F = \|Q^* G \Delta^* Q\|_F = \|G\|_F = \pi_2(A).$$

The same arguments yield  $\hat{\pi}_1(A) = \hat{\pi}_2(A)$   $\square$

Lemma 2.  $\pi_1(A) \leq \pi_3(A) \leq \pi_1(A)(2 + \pi_1(A))$ ,  $\hat{\pi}_1(A) \leq \hat{\pi}_3(A) \leq \hat{\pi}_1(A)(2 + \hat{\pi}_1(A))$ .

Proof. Assume  $A = U \Sigma V^*$ , the singular value decomposition of  $A$  and let be  $\Sigma = I + D$ . Then  $D_{ii} \geq -1$  and

$$\|A - U V^*\|_F = \|U D V^*\|_F = \|D\|_F \geq \pi_1(A).$$

Furthermore

$$\pi_3(A) = \|A^* A - I\|_F = \|V(I+D)^2 V^* - I\|_F = \|2D + D^2\|_F \geq \|D\|_F \geq \pi_1(A),$$

for  $D_{ii} \geq -1$ . With  $\pi_1(A) = \|A - V\|_F$ ,  $V \in U$  and  $A = V + E$  one finds

$$\begin{aligned} \pi_3(A) &= \|A^* A - I\|_F = \|(V+E)^*(V+E) - I\|_F = \|V^* E + E^* V + E^* E\|_F \\ &\leq 2 \|E\|_F + \|E\|_F^2 = 2 \pi_1(A) + \pi_1^2(A). \end{aligned}$$

The same arguments yield  $\hat{\pi}_1(A) \leq \hat{\pi}_3(A) \leq \hat{\pi}_1(A)(2 + \hat{\pi}_1(A))$   $\square$

For the proof of (C15) let be  $A = U \Sigma V^*$  a singular value decomposition of  $A$ . Then  $H_1 = U \Sigma U^*$  and  $H_2 = V \Sigma V^*$ . Then, with  $W = U^* V$  one has

$$U^* (H_1 - H_2) V = \Sigma W - W \Sigma.$$

Without loss of generality we may assume that equal singular values are consecutive in the diagonal of  $\Sigma$ . When the multiplicities of the  $r$  ( $2 \leq r \leq n$ ) distinct singular values  $\sigma_{k_1}, \dots, \sigma_{k_r}$  are  $n_1, \dots, n_r$  resp. then we write

$$\Sigma = \begin{pmatrix} \sigma_{k_1} I_1 & 0 & & 0 \\ 0 & \sigma_{k_2} I_2 & \dots & 0 \\ \vdots & \vdots & & \vdots \\ 0 & 0 & \dots & \sigma_{k_r} I_r \end{pmatrix}, \quad I_j \in C^{n_j, n_j},$$

with  $\sigma_{k_i} \neq \sigma_{k_j}$  for  $i \neq j$ . Hence  $\tau = \min_{i \neq j} |\sigma_{k_i} - \sigma_{k_j}|$ . Similarly we partition  $W$ . So we obtain

$$(4.16) \quad \mu_5^2 = \|U^*(H_1 - H_2)U\|_F^2 = \sum_{i \neq j} (\sigma_{k_i} - \sigma_{k_j})^2 \|W_{ij}\|_F^2 \geq \tau^2 \sum_{i \neq j} \|W_{ij}\|_F^2,$$

where  $W_{ij} \in C^{n_i, n_j}$  is a block in the partitioned matrix  $W$ . (4.16) yields

$$(4.17) \quad \sum_{i \neq j} \|W_{ij}\|_F^2 \leq (\mu_5/\tau)^2 =: \epsilon^2 < 1.$$

In each diagonal block  $W_{jj} \in C^{n_j, n_j}$  of  $W$  the length of each column is in the interval  $[(1-\epsilon^2)^{\frac{1}{2}}, 1]$  and the inner product of each pair of columns of each  $W_{jj}$  is smaller than  $\epsilon^2$ . Hence

$$(4.18) \quad \pi_3(W_{jj}) = \|W_{jj}^* W_{jj} - I\|_F \leq n_j \epsilon, \quad j = 1, \dots, r.$$

In view of lemma 2 there exists a unitary  $Q_j \in C^{n_j, n_j}$  and a unitary diagonal  $\Delta_j \in C^{n_j, n_j}$  such that

$$\pi_1(W_{jj}) = \pi_2(W_{jj}) = \|Q_j^* W_{jj} Q_j \Delta_j - I\|_F \leq \pi_3(W_{jj}) \leq n_j \epsilon, \\ j = 1, \dots, r.$$

These matrices  $Q_j$  and  $\Delta_j$  are used in the formation of the block diagonal matrices

$$D_1 = \begin{pmatrix} Q_1 & 0 & \dots & 0 \\ 0 & Q_2 & \dots & 0 \\ \vdots & \vdots & & \vdots \\ 0 & 0 & \dots & Q_r \end{pmatrix}, \quad D_2 = \begin{pmatrix} Q_1 \Delta_1 & 0 & \dots & 0 \\ 0 & Q_2 \Delta_2 & \dots & 0 \\ \vdots & \vdots & & \vdots \\ 0 & 0 & \dots & Q_r \Delta_r \end{pmatrix}.$$

Let be

$$\hat{U} := U D_1, \quad \hat{V} := V D_2.$$

Then

$$\hat{U}^* A \hat{V} = D_1^* \Sigma D_2 = \text{diag}(\sigma_{k_j} \Delta_j) \in D$$

and

$$\begin{aligned} \|\hat{V} - \hat{U}\|_F^2 &= \|\hat{U}^* \hat{V} - I\|_F^2 = \|D_1^* U^* V D_2 - I\|_F^2 \\ &= \|\text{diag}(Q_j^* W_{jj} Q_j \Delta_j - I)\|_F^2 + \sum_{i \neq j} \|W_{ij}\|^2. \end{aligned}$$

With (4.17) one has

$$\|\hat{V} - \hat{U}\|_F^2 \leq \sum_{j=1}^r n_j^2 \varepsilon^2 + \varepsilon^2 = \varepsilon^2 (1 + \sum_{j=1}^r n_j^2) \leq \varepsilon^2 (1 + n^2).$$

This means

$$\mu_8 \leq \varepsilon \sqrt{1+n^2} \leq (n+1) \mu_5 / \tau.$$

For the proof of (C16), as can be done without loss of generality, we assume that  $M := (A + A^*)/2 = \text{diag}(\alpha_j)$ .

When the multiplicities of the  $r$  distinct eigenvalues

$\alpha_{k_1}, \dots, \alpha_{k_r}$  ( $1 \leq r \leq n$ ) are  $n_1, \dots, n_r$  resp. we write

$$M = \frac{A+A^*}{2} = \begin{pmatrix} \alpha_{k_1} I_1 & 0 & \dots & 0 \\ 0 & \alpha_{k_2} I_2 & \dots & 0 \\ \vdots & \vdots & & \vdots \\ 0 & 0 & \dots & \alpha_{k_r} I_r \end{pmatrix}, \quad I_j \quad R^{n_j, n_j}.$$

The matrix  $G = (A-A^*)/(2i)$  is partitioned in the same way and we assume that each diagonal block  $G_{jj}$  has been unitarily diagonalized. Hence

$$G = \frac{A-A^*}{2} = \text{diag}(\gamma_1, \dots, \gamma_n) + \begin{pmatrix} 0 & G_{12} & \dots & G_{1n} \\ G_{21} & 0 & \dots & G_{2n} \\ \vdots & \vdots & & \vdots \\ G_{n1} & G_{n2} & \dots & 0 \end{pmatrix}.$$

Then

$$(4.19) \quad \frac{1}{4} \|A^*A - AA^*\|_F^2 = \|GM - MG\|_F^2 = \sum_{i \neq j} (\alpha_{k_i} - \alpha_{k_j})^2 \|G_{ij}\|_F^2 \geq \delta^2 \sum_{i \neq j} \|G_{ij}\|_F^2,$$

$$\text{where } \delta = \min_{i \neq j} |\alpha_{k_i} - \alpha_{k_j}| = \min_{\alpha_i \neq \alpha_j} |\alpha_i - \alpha_j|.$$

Hence (4.19) implies

$$(4.20) \quad \left( \sum_{i \neq j} \|G_{ij}\|_F^2 \right)^{\frac{1}{2}} \leq \frac{1}{2} \mu_2^2 / \delta.$$

With the theorem of Hoffman-Wielandt [8] one derives that there exists a permutation  $p$  of the eigenvalues  $\beta_j$  of  $G$  such that

$$\left( \sum_{j=1}^n (\beta_{p(j)} - \gamma_j)^2 \right)^{\frac{1}{2}} \leq \frac{1}{2} \mu_2^2 / \delta.$$

$$\text{So } |\beta_{p(j)} - \gamma_j| \leq \frac{1}{2} \mu_2^2 / \delta, \quad j = 1, \dots, n, \text{ and}$$

$$(4.21) \quad |\gamma_i - \gamma_j| \geq |\beta_{p(i)} - \beta_{p(j)}| - |\beta_{p(i)} - \gamma_i| - |\beta_{p(j)} - \gamma_j| \geq \tilde{\delta} - \mu_2^2/\delta, \\ i \neq j,$$

where

$$\tilde{\delta} = \min_{i \neq j} |\beta_i - \beta_j|,$$

the minimal distance of the distinct eigenvalues of  $G$ .

Now we investigate the distance between  $e_1 = (1, 0, \dots, 0)^T$  and the normalized eigenvalues of  $G$  corresponding with  $\beta_{p(1)}$ . For that purpose  $G$  is partitioned in the following way

$$G = \begin{pmatrix} \gamma_1 & \vdots & 0^T \\ \vdots & \ddots & \vdots \\ 0 & \vdots & \text{diag}(\gamma_2, \dots, \gamma_n) \end{pmatrix} + \begin{pmatrix} 0 & \vdots & g^T \\ \vdots & \ddots & \vdots \\ g & \vdots & \tilde{G} \end{pmatrix}, \quad \tilde{G} \in \mathbb{C}^{n-1, n-1}.$$

Since  $\min_{2 \leq i \leq n} |\gamma_1 - \gamma_i| \geq \tilde{\delta} - \mu_2^2/\delta$  as follows from (3.21) and  $\|\tilde{G}\|_F \leq \frac{1}{2} \mu_2^2/\delta$  as follows from (4.20),

$$\min_{2 \leq i \leq n} |\gamma_1 - \gamma_i| - \|\tilde{G}\|_F \geq \tilde{\delta} - \frac{3}{2} \mu_2^2/\delta > 0.$$

Moreover,  $\|g\|_2 \leq \frac{1}{2} \mu_2^2/\delta$ , once again as a consequence of (4.20). Applying Stewart's theorem on the sensitivity of an invariant subspace [7] we conclude that there exists a vector  $p_1 \in \mathbb{C}^{n-1}$  satisfying

$$\|p_1\|_2 \leq 2 \|g\|_2 / (\tilde{\delta} - \frac{3}{2} \mu_2^2/\delta) \leq \mu_2^2 / (\delta \tilde{\delta} - \frac{3}{2} \mu_2^2)$$

such that  $(1, p_1^T)^T \in \mathbb{C}^n$  is an eigenvector of  $G$ .

The distance between  $e_1$  and  $v_1 := (1 + \|p_1\|^2)^{-\frac{1}{2}} (1, p_1^T)^T$  equals

$$(2(1 - (1 + \|p_1\|^2)^{-\frac{1}{2}}))^{\frac{1}{2}} \leq \|p_1\|_2 \leq \mu_2^2 / (\delta \tilde{\delta} - \frac{3}{2} \mu_2^2).$$

By assumption

$$\mu_2^2 / (\delta\hat{\delta} - \frac{3}{2} \mu_2^2) < \frac{1}{2} \sqrt{2}$$

so  $\|v_1 - e_j\| > \|v_1 - e_1\|$  for each unit vector  $e_j$  with  $j > 1$ .

In an analogous manner one can derive the existence of eigenvalues  $v_j$ ,  $j = 2, \dots, w$ , such that  $\|v_j - e_j\| < \mu_2^2 / (\delta\hat{\delta} - \frac{3}{2} \mu_2^2) < \frac{1}{2} \sqrt{2}$ .

Let be  $v_1, \dots, v_n$  the columns of  $V \in U$ . Then

$$\mu_7 \leq \|V - I\|_F \leq \sqrt{n} \mu_2^2 (\delta\hat{\delta} - \frac{3}{2} \mu_2^2)^{-1}.$$

This proves inequality (C16) and ends the proof of theorem 2.  $\square$



### References

1. P.J. Eberlein, On measures of non-normality for matrices, Amer. Math. Monthly, 72 (1965), pp. 995-996.
2. P. Henrici, Bounds for iterates, inverses, spectral variation and fields of values of nonnormal matrices, Numer. Math., 4 (1962), pp. 24-40.
3. R. Kress, H.L. de Vries, R. Wegemann, On nonnormal matrices, Lin. Alg. Appl., 8, (1974), pp. 109-120.
4. M. Marcus - H. Minc, A survey of matrix theory and matrix inequalities, Allyn and Bacon, 1964.
5. A. Ruhe, On the closeness of eigenvalues and singular values for almost normal matrices, Lin. Alg. Appl., 11 (1975), pp. 87-94.
6. R.A. Smith, The condition number of the matrix eigenvalue problem, Numer. Math., 10 (1967), pp. 232-240.
7. G.W. Stewart, Error and Perturbation Bounds for Subspaces associated with certain Eigenvalue Problems, Siam Review, 15 (1973), pp. 727-764.
8. A.J. Hoffman, H.W. Wielandt, The variation of the spectrum of a normal matrix, Duke Math. J., 20, (1953), pp. 37-39.

IN 1983 REEDS VERSCHENEN

- 126 H.H. Tigelaar  
Identification of noisy linear systems with multiple arma inputs.
- 127 J.P.C. Kleijnen  
Statistical Analysis of Steady-State Simulations: Survey of Recent Progress.
- 128 A.J. de Zeeuw  
Two notes on Nash and Information.
- 129 H.L. Theuns en A.M.L. Passier-Grootjans  
Toeristische ontwikkeling - voorwaarden en systematiek; een selectief literatuuroverzicht.
- 130 J. Plasmans en V. Somers  
A Maximum Likelihood Estimation Method of a Three Market Disequilibrium Model.
- 131 R. van Montfort, R. Schippers, R. Heuts  
Johnson  $S_U$ -transformations for parameter estimation in arma-models when data are non-gaussian.
- 132 J. Glombowski en M. Krüger  
On the Rôle of Distribution in Different Theories of Cyclical Growth.
- 133 J.W.A. Vingerhoets en H.J.A. Coppens  
Internationale Grondstoffenovereenkomsten.  
Effecten, kosten en oligopolisten.
- 134 W.J. Oomens  
The economic interpretation of the advertising effect of Lydia Pinkham.
- 135 J.P.C. Kleijnen  
Regression analysis: assumptions, alternatives, applications.
- 136 J.P.C. Kleijnen  
On the interpretation of variables.
- 137 G. van der Laan en A.J.J. Talman  
Simplicial approximation of solutions to the nonlinear complementarity problem with lower and upper bounds.

IN 1984 REEDS VERSCHENEN

- 138 G.J. Cuypers, J.P.C. Kleijnen en J.W.M. van Rooyen  
Testing the Mean of an Asymmetric Population:  
Four Procedures Evaluated
- 139 T. Wansbeek en A. Kapteyn  
Estimation in a linear model with serially correlated errors when  
observations are missing
- 140 A. Kapteyn, S. van de Geer, H. van de Stadt, T. Wansbeek  
Interdependent preferences: an econometric analysis
- 141 W.J.H. van Groenendaal  
Discrete and continuous univariate modelling
- 142 J.P.C. Kleijnen, P. Cremers, F. van Belle  
The power of weighted and ordinary least squares with estimated  
unequal variances in experimental design
- 143 J.P.C. Kleijnen  
Superefficient estimation of power functions in simulation  
experiments
- 144 P.A. Bekker, D.S.G. Pollock  
Identification of linear stochastic models with covariance  
restrictions.
- 145 Max D. Merbis, Aart J. de Zeeuw  
From structural form to state-space form
- 146 T.M. Doup and A.J.J. Talman  
A new variable dimension simplicial algorithm to find equilibria on  
the product space of unit simplices.
- 147 G. van der Laan, A.J.J. Talman and L. Van der Heyden  
Variable dimension algorithms for improper labellings.
- 148 G.J.C.Th. van Schijndel  
Dynamic firm behaviour and financial leverage clienteles
- 149 M. Plattel, J. Peil  
The ethico-political and theoretical reconstruction of contemporary  
economic doctrines
- 150 F.J.A.M. Hoes, C.W. Vroom  
Japanese Business Policy: The Cash Flow Triangle  
an exercise in sociological demystification
- 151 T.M. Doup, G. van der Laan and A.J.J. Talman  
The  $(2^{n+1}-2)$ -ray algorithm: a new simplicial algorithm to compute  
economic equilibria

IN 1984 REEDS VERSCHENEN (vervolg)

- 152 A.L. Hempenius, P.G.H. Mulder  
Total Mortality Analysis of the Rotterdam Sample of the Kaunas-Rotterdam Intervention Study (KRIS)
- 153 A. Kapteyn, P. Kooreman  
A disaggregated analysis of the allocation of time within the household.
- 154 T. Wansbeek, A. Kapteyn  
Statistically and Computationally Efficient Estimation of the Gravity Model.
- 155 P.F.P.M. Nederstigt  
Over de kosten per ziekenhuisopname en levensduurmodellen
- 156 B.R. Meijboom  
An input-output like corporate model including multiple technologies and make-or-buy decisions
- 157 P. Kooreman, A. Kapteyn  
Estimation of Rationed and Unrationed Household Labor Supply Functions Using Flexible Functional Forms
- 158 R. Heuts, J. van Lieshout  
An implementation of an inventory model with stochastic lead time
- 159 P.A. Bekker  
Comment on: Identification in the Linear Errors in Variables Model
- 160 P. Meys  
Functies en vormen van de burgerlijke staat  
Over parlementarisme, corporatisme en autoritair etatisme
- 161 J.P.C. Kleijnen, H.M.M.T. Denis, R.M.G. Kerckhoffs  
Efficient estimation of power functions
- 162 H.L. Theuns  
The emergence of research on third world tourism: 1945 to 1970;  
An introductory essay cum bibliography
- 163 F. Boekema, L. Verhoef  
De "Grijze" sector zwart op wit  
Werklozenprojecten en ondersteunende instanties in Nederland in kaart gebracht
- 164 G. van der Laan, A.J.J. Talman, L. Van der Heyden  
Shortest paths for simplicial algorithms
- 165 J.H.F. Schilderincx  
Interregional structure of the European Community  
Part II: Interregional input-output tables of the European Community 1959, 1965, 1970 and 1975.

- 166 P.J.F.G. Meulendijks  
An exercise in welfare economics (I)





Bibliotheek K. U. Brabant



17 000 01059776 4